



Sepsis Fact Sheet

Definition: Sepsis is the body's overwhelming and life-threatening response to infection, which can lead to tissue damage, organ failure, and death.

Who it Hurts: While sepsis is an equal-opportunity killer, impacting the sick, the well, and people of all ages, some groups are more likely to be affected. These include very young children, older adults, and those with a weakened immune system.

Prevention: The risk of sepsis can be reduced by preventing or quickly identifying and managing infections. This includes practicing good hygiene, staying current with vaccinations, and seeking treatment when infections are suspected.

Treatment: Sepsis is a medical emergency that requires urgent attention and rapid treatment for survival. Sepsis can be treated and, in many instances, lives are saved by using existing and proven protocols.

Recovery: Many individuals fully recover from sepsis, while others may have long-lasting effects, such as amputations or organ dysfunction, like kidney failure. Other aftereffects of sepsis are less obvious, such as memory loss, anxiety, or depression.

Symptoms: When it comes to sepsis, remember **It's About TIME™**:

T – Temperature – higher or lower than normal

I – Infection – may have signs or symptoms of infection

M – Mental Decline – confused, sleepy, difficult to rouse

E – Extremely ill – severe pain, discomfort, shortness of breath

If you **suspect sepsis** (observe a combination of these symptoms), see your medical professional immediately, CALL 911, or go to a hospital with an advocate and say, **“I AM CONCERNED ABOUT SEPSIS.”**

Critical Facts:

- Sepsis is the leading cause of death in U.S. hospitals.¹
- Approximately 6% of all hospitalizations are due to sepsis and 35% of all deaths in-hospital are due to sepsis.²
- Sepsis is the leading cause of readmissions to the hospital, with as many as 19% of people originally hospitalized with sepsis re-hospitalized within 30 days and about 40% rehospitalized within 90 days.^{3, 4, 5}
- As many as 87% of sepsis cases originate in the community and not in the hospital.²
- Mortality from sepsis increases by as much as 8% for every hour treatment is delayed.^{6, 7, 8} As many as 80% of sepsis deaths could be prevented with rapid diagnosis and treatment.⁶
- Sepsis is the most common complication observed in severe cases of COVID-19.^{9, 10}

Human Cost:

- Sepsis affects an estimated 49 million people worldwide each year, including more than 20 million children under age 5, and nearly 5 million older children and adolescents (ages 5-19).¹¹
- Sepsis takes 11 million lives around the world each year, contributing to 20% of all deaths globally and taking more lives than cancer.^{11, 12} This is more than 20 deaths every minute.¹³
- More than 1.7 million people in the U.S. are diagnosed with sepsis each year – one every 20 seconds – and the incidence is rising.^{2, 14, 15}
- 270,000 people die from sepsis every year in the U.S. – one every 2 minutes; more than from prostate cancer, breast cancer and opioid overdose combined.^{2, 16, 17}
- It is estimated that each year, more than 80,000 sepsis survivors are discharged to hospice after a sepsis hospitalization.²
- Each year, more than 75,000 children in the U.S. develop severe sepsis and 6,800 of these children die, more than from pediatric cancers.^{18, 19}
- Sepsis causes at least 261,000 maternal deaths every year worldwide and is driving increases in pregnancy-related deaths in the U.S.^{20, 21}
- In 2012, there were more than 13,700 sepsis-related amputations in the U.S. This works out to an average of 38 amputations per day.²² Approximately 1% of sepsis survivors undergo one or more surgical amputations of a limb or digit as a result of sepsis.²³
- Sepsis survivors have a shortened life expectancy, are more likely to suffer from an impaired quality of life, and often experience worsened cognitive (mental) and physical function.^{5, 15, 24, 25, 26}
- Missed infections are the third most common cause of diagnostic errors identified in medical malpractice cases. Sepsis is the most common condition among missed infections in diagnostic errors.²⁷

Economic Cost:

- Sepsis is the #1 cost of hospitalization in the U.S.²⁸ Costs for acute sepsis hospitalization and skilled nursing are estimated to be \$62 billion annually.²⁹ This is only a portion of all sepsis-related costs, since there are substantial additional costs after discharge for many.
- The average cost per hospital stay for sepsis is double the average cost per stay across all other conditions.³⁰
- Sepsis is the #1 cause of readmission to the hospital, costing more than \$3.5 billion each year.^{4, 28}

Awareness:

- More than 34% of U.S. adults have NEVER heard of sepsis and just 14% can name the common symptoms.³¹

To find out more please visit [Sepsis.org](https://www.sepsis.org)

Sources:

1. Liu V, et al. JAMA. 2014;312(1):90-92.
<http://jama.jamanetwork.com/article.aspx?articleid=1873131&resultClick=3>
2. Rhee C, et al. JAMA. 2017;318(13):1241-1249. <http://jamanetwork.com/journals/jama/fullarticle/2654187>
3. Fingar K, Washington R. HCUP Statistical Brief #196. November 2015. Agency for Healthcare Research and Quality, Rockville, MD. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb196-Readmissions-Trends-High-Volume-Conditions.jsp>
4. Gadre SK, et al. Chest. 2019;155(3):483-490. [https://journal.chestnet.org/article/S0012-3692\(18\)32895-2/abstract](https://journal.chestnet.org/article/S0012-3692(18)32895-2/abstract)
5. Prescott H and Angus D. JAMA. 2018;319(1):91. <https://jamanetwork.com/journals/jama/fullarticle/2667724>
6. Kumar A, et al. Crit Care Med. 2006;34(6):1589-1596. <https://pubmed.ncbi.nlm.nih.gov/16625125/>
7. Seymour CW, et al. N Engl J Med. 2017;376(23):2235-2244.
<https://www.nejm.org/doi/full/10.1056/NEJMoa1703058>
8. Liu VX, et al. Am J Respir Crit Care Med. 2017;196(7):856-863.
<https://www.atsjournals.org/doi/full/10.1164/rccm.201609-1848OC>
9. Zhou F, et al. Lancet. 2020; 395 (10229), 1054-1062.
[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)30566-3/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30566-3/fulltext)
10. Chen T, et al. BMJ. 2020; 368:m1091. <https://www.bmj.com/content/368/bmj.m1091>
11. Rudd KE, et al. Lancet. 2020;395(10219):200-211
[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(19\)32989-7/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(19)32989-7/fulltext)
12. Cancer, World Health Organization, 12 September 2018. Accessed June 2, 2020. https://www.who.int/health-topics/cancer#tab=tab_1
13. Kempker JA, Martin GS. Lancet. 2020;395(10219):168-170.
[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(19\)33065-X/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(19)33065-X/fulltext)
14. Hall, MJ, et al. NCHS Data Brief, No 62. Hyattsville, MD: National Center for Health Statistics. 2011.
<https://www.cdc.gov/nchs/data/databriefs/db62.pdf>
15. Buchman TG, et al. Sepsis Among Medicare Beneficiaries: 1. The Burdens of Sepsis, 2012-2018. Crit Care Med. 2020;48(3):276-288.
https://journals.lww.com/ccmjournal/fulltext/2020/03000/sepsis_among_medicare_beneficiaries__1__the.2.aspx
16. Cancer Stat Facts: Cancer of Any Site, National Cancer Institute. Accessed June 2, 2020.
<https://seer.cancer.gov/statfacts/html/all.html>
17. Overdose Death Rates, National Institute on Drug Abuse, March 2020. Accessed June 2, 2020.
<https://www.drugabuse.gov/related-topics/trends-statistics/overdose-death-rates>
18. Hartman ME, et al. Pediatr Crit Care Med. 2013;14(7):686-693.
<https://www.ncbi.nlm.nih.gov/pubmed/23897242>
19. Cancer in Children and Adolescents. National Cancer Institute. Retrieved 3/31/20.
<https://www.cancer.gov/types/childhood-cancers/child-adolescent-cancers-fact-sheet>
20. Say L, et al. Lancet Glob Health 2014;2(6):e323-333.
<http://www.sciencedirect.com/science/article/pii/S2214109X1470227X>
21. Pregnancy Mortality Surveillance System, CDC, February 4, 2020.
<http://www.cdc.gov/reproductivehealth/maternalinfanthealth/pmss.html>
22. Healthcare Cost and Utilization Project, Nationwide Inpatient Sample, 2012. Accessed April 6, 2016
23. Reitz K, et al. Am J Resp Crit Care Med, AJRCCM Conference Abstracts 2019, 201: A7153.
https://www.atsjournals.org/doi/abs/10.1164/ajrccm-conference.2019.199.1_MeetingAbstracts.A7153

24. Winters BD, et al. *Critical Care Medicine Journal*. 2010;38(5):8.2010.
<http://www.ncbi.nlm.nih.gov/pubmed/20308885>
25. Lund-Sorensen H, et al. 2016 <https://jamanetwork.com/journals/jamapsychiatry/article-abstract/2542681>
26. Iwashyna TJ, et al. *JAMA*. 2010;304(16):8. <https://jamanetwork.com/journals/jama/fullarticle/186769>
27. Newman-Toker DE, et al. *Diagnosis*. 2019;6(3):227-240. <https://pubmed.ncbi.nlm.nih.gov/31535832/>
28. Torio C, Moore B. HCUP Statistical Brief #204. May 2016. Agency for Healthcare Research and Quality, Rockville, MD. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb204-Most-Expensive-Hospital-Conditions.pdf>
29. Buchman TG, Simpson SQ, Sciarretta KL, et al. Sepsis Among Medicare Beneficiaries: 3. The Methods, Models, and Forecasts of Sepsis, 2012-2018. *Crit Care Med*. 2020;48(3):302-318.
https://journals.lww.com/ccmjournal/FullText/2020/03000/Sepsis_Among_Medicare_Beneficiaries_3_The.4.aspx
30. Pfunter A, et al. HCUP Statistical Brief #168. December 2013. Agency for Healthcare Research and Quality, Rockville, MD. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb168-Hospital-Costs-United-States-2011.jsp>
31. 2019 Sepsis Awareness Survey Results, Sepsis Alliance. <http://www.sepsis.org/2019-sepsis-awareness-survey/>

Revised 6/2020