



Sepsis and Health Equity Fact Sheet*

Introduction

In the United States, differences in health and mortality are influenced by both socio-economic status (SES) and race/ethnicity.¹ Sepsis, the body's overwhelming and life-threatening response to infection, is no exception.

Sepsis is the leading cause of death in U.S. hospitals,² and the #1 cost of hospitalization at \$53 billion annually.^{3, 4} More than 1.7 million people in the U.S. are diagnosed with sepsis each year, with an estimated 270,000 deaths every year in the U.S. alone; more than from prostate cancer, breast cancer, and opioid overdose combined.^{5, 6, 7, 8} Those who survive sepsis have a shortened life expectancy, are more likely to suffer from an impaired quality of life, and often experience worsened mental and physical function.⁹

10, 11, 12, 13

Many factors link SES and race/ethnicity to health disparities: racism; poverty; differential access to resources that promote health; unequal access to quality healthcare and treatments; different rates and types of stressors; healthiness of one's residential and neighborhood environment; and the development of preventable chronic health conditions.^{14, 15}

For Sepsis Alliance, health equity means eliminating preventable disparities in sepsis incidence, morbidity, and mortality. This fact sheet looks at what is currently known about these disparities, as well as disparities in sepsis awareness and treatment.

Research on sepsis disparities to date has focused primarily on socioeconomic status and/or comparisons between Blacks or African Americans, Hispanics, and whites. More research is also needed on smaller minority populations, such as Native Americans, Pacific Islanders, and Asian Americans, who are often left out of health studies due to low numbers, or combined with other groups within combined racial/ethnic group designations such as "Other" or "Asian and Pacific Islander."

Racial/Ethnic Disparities in the Incidence and Mortality of Sepsis

- Black and "other nonwhite" individuals have nearly twice the incidence of sepsis as whites (1.89 times the risk for Blacks, and 1.9 times the risk for "other nonwhite" individuals).¹⁶

- Black and Hispanic individuals also have a higher incidence of severe sepsis as compared to whites (1.7 times the rate for Blacks, and 1.1 times the rate for Hispanics).¹⁷
- Blacks bear nearly twice the burden of sepsis deaths relative to the size of the Black population, as compared to whites.^{18, 19} Sepsis deaths among American Indians/Alaskan Natives and Hispanics are also elevated as compared to whites (1.24 times the risk for American Indians/Alaskan Natives and 1.14 times the risk for Hispanics).²⁰
- American Indians and Alaskan Natives in the Indian Health Service area are 1.6 times more likely to die from sepsis as the national average.⁴⁰
- Research from Hawaii finds that Native Hawaiians have almost twice the burden of sepsis deaths as compared to whites.²¹

Racial/Ethnic Disparities in Patient Care and Outcomes

- Black patients admitted to the emergency room are assigned to significantly lower priority status and experience significantly longer wait times (10.9 minutes longer on average) as compared to case-matched white patients.²²
- Non-Hispanic Black children admitted to the emergency room are less likely to be treated for sepsis than non-Hispanic white children.²³
- Sepsis deaths among prostate cancer patients are higher for racial and ethnic minorities. Blacks with prostate cancer have 1.9 times the risk, American-Indians 2.1 times the risk, and Asian and Pacific Islanders 1.9 times the risk of sepsis death as compared to whites with prostate cancer.²⁴
- Black children are 30% more likely than white children to develop sepsis after surgery.²⁵
- Research indicates more than twice the risk of severe maternal sepsis for maternal patients who are Black as compared to white maternal patients.²⁶
- Preliminary research in a rural emergency department serving a Native American population found that 27.9% of admissions and transfers met sepsis criteria, higher than the 6% average for U.S. hospitals.^{27, 8}
- Black patients with severe sepsis have 8% higher case fatality rates than white

sepsis patients, are 4% less likely to be admitted to the ICU, and are 9% more likely to die if admitted to the ICU.¹⁷

- Other research also indicates that mortality among hospitalized sepsis patients at any level of severity is significantly higher for racial and ethnic minorities compared to white patients. Hospitalized Black or Hispanic sepsis patients are 7% more likely to die than white, while those categorized as "Asian and Pacific Islander" or "Other" race are 18% or 21% more likely to die than white patients, respectively, after taking other patient characteristics into account.²⁸
- Research conducted at a diverse urban medical center found that Asian sepsis patients were 57% more likely to die than their white counterparts.²⁹
- Children with severe sepsis or septic shock who are Black or Hispanic are approximately 25% more likely to die than non-Hispanic white children.³⁰
- Limited English proficiency is associated with an 80% higher mortality risk among sepsis patients.²⁹
- In comparing Black vs. white patients, racial differences in sepsis are due to both a 39% higher rate of infection and a 29% higher rate of organ dysfunction complications in those infections.¹⁹
- Black and Native American patients are more likely to be readmitted following a sepsis hospitalization as compared to their white counterparts (1.29 times the risk for Blacks and 2.39 times the risk for Native Americans).³¹

Poverty and Socioeconomic Status

Adults with lower levels of education, income, and/or material resources, are at greater risk of sepsis mortality.¹⁸ For example,

- Adults without a high school diploma have over 2.5 times the risk of dying from sepsis as those with a doctorate;
- Adults without a telephone number in the home are 1.6 times as likely to die of sepsis as those with; and
- Adults below the poverty line have over three to four times the risk of dying of sepsis as compared to adults whose family income is at least five times the poverty line.¹⁸

- Adult patients without health insurance are more likely to die of sepsis than privately insured patients, and are less than half as likely to be discharged to a nonhospital healthcare facility or discharged with home healthcare.³²

Infants and children with lower levels of income and access to insurance are at greater risk of sepsis mortality.^{33,34}

- Infants from lower income families are 20% more likely to die from sepsis.³³
- Infants from families without health insurance are 3 times more likely to die from sepsis.³³
- Children with severe sepsis or septic shock with public insurance are more likely to die than children with private or other types of insurance.³⁴

Access to intensive care and hospitals with lower mortality rates, is related to income and minority status.

- Health care resources, including the availability of ICU beds, are more plentiful in wealthier communities. In the U.S., nearly half (49%) of the lowest income communities have no ICU beds, whereas only 3% of the highest income communities have no ICU beds.³⁵
- Sepsis patients of any race/ethnicity who are treated in predominantly minority-serving hospitals have higher rates of in-hospital mortality.^{36, 37, 38}

Awareness and Knowledge

- Sepsis awareness is significantly lower for Blacks than for whites. In a recent survey conducted by Sepsis Alliance, only 49% of respondents identifying as Black had heard the term sepsis, as compared to 76% of white-identifying respondents.³⁹
- Only 5% of Black respondents identified all four common symptoms of sepsis included in Sepsis Alliance's annual survey, as compared to 18% of white survey respondents.³⁹
- White survey respondents were significantly more likely to know someone

who had sepsis or had it themselves (37%), as compared to Black respondents (26%).³⁹

- Income differences were also significantly related to sepsis awareness, with 15% more of the higher income respondents having heard the term 'sepsis' as compared to the lower income respondents.³⁹

* The Sepsis and Health Equity Fact Sheet was made possible in part with an unrestricted grant from Roche Diagnostics.

References

1. Phelan, J. C., & Link, B. G. (2015). Is racism a fundamental cause of inequalities in health?. *Annual Review of Sociology*, 41, 311-330.
2. Liu, V., Escobar, G. J., Greene, J. D., Soule, J., Whippy, A., Angus, D. C., & Iwashyna, T. J. (2014). Hospital deaths in patients with sepsis from 2 independent cohorts. *JAMA*, 312(1), 90-92.
3. Buchman, T. G., Simpson, S. Q., Sciarretta, K. L., Finne, K. P., Sowers, N., Collier, M., ... & Wax, M. (2020b). Sepsis Among Medicare Beneficiaries: 3. The Methods, Models, and Forecasts of Sepsis, 2012–2018. *Critical Care Medicine*, 48(3), 302.
4. Torio, C. M., & Moore, B. J. (2016). National inpatient hospital costs: the most expensive conditions by payer, 2013: statistical brief# 204. *Healthcare cost and utilization project (HCUP) statistical briefs*, 2006-2016.
5. Cancer Stat Facts: Cancer of Any Site, National Cancer Institute. (n.d.) Retrieved June 2, 2020. <https://seer.cancer.gov/statfacts/html/all.html>
6. Overdose Death Rates, National Institute on Drug Abuse. (2020, March) <https://www.drugabuse.gov/related-topics/trends-statistics/overdose-death-rates>
7. Rudd, K. E., Johnson, S. C., Agesa, K. M., Shackelford, K. A., Tsoi, D., Kievlan, D. R., ... & Fleischmann-Struzek, C. (2020). Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *The Lancet*, 395(10219), 200-211.
8. Rhee, C., Dantes, R., Epstein, L., Murphy, D. J., Seymour, C. W., Iwashyna, T. J., ... & Jernigan, J. A. (2017). Incidence and trends of sepsis in US hospitals using clinical vs claims data, 2009-2014. *JAMA*, 318(13), 1241-1249.
9. Buchman, T. G., Simpson, S. Q., Sciarretta, K. L., Finne, K. P., Sowers, N., Collier, M., ... & Wax, M. (2020a). Sepsis Among Medicare Beneficiaries: 1. The Burdens of Sepsis, 2012–2018. *Critical Care Medicine*, 48(3), 276.
10. Iwashyna, T. J., Ely, E. W., Smith, D. M., & Langa, K. M. (2010). Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA*, 304(16), 1787-1794.
11. Lund-Sørensen, H., Benros, M. E., Madsen, T., Sørensen, H. J., Eaton, W. W., Postolache, T. T., ... & Erlangsen, A. (2016). A nationwide cohort study of the

association between hospitalization with infection and risk of death by suicide. *JAMA psychiatry*, 73(9), 912-919.

12. Prescott, H. C., & Angus, D. C. (2018). Postsepsis morbidity. *JAMA*, 319(1), 91-91.

13. Winters, B. D., Eberlein, M., Leung, J., Needham, D. M., Pronovost, P. J., & Sevransky, J. E. (2010). Long-term mortality and quality of life in sepsis: a systematic review. *Critical Care Medicine*, 38(5), 1276-1283.

14. DiMeglio, M., Dubensky, J., Schadt, S., Potdar, R. and Laudanski, K. (2018) Factors Underlying Racial Disparities in Sepsis Management, *Healthcare*, 6(4), 133.

15. Vogel, T. R. (2012) Update and review of racial disparities in sepsis, *Surgical Infections*, 13(4), 203-208.

16. Martin, G. S., Mannino, D. M., Eaton, S. and Moss, M. (2003) The epidemiology of sepsis in the United States from 1979 through 2000, *New England Journal of Medicine*, 348(16), 1546-1554.

17. Barnato, A. E., Alexander, S. L., Linde-Zwirble, W. T. and Angus, D. C. (2008) Racial variation in the incidence, care, and outcomes of severe sepsis: analysis of population, patient, and hospital characteristics, *American Journal of Respiratory and Critical Care Medicine*, 177(3), 279-284.

18. Kempker, J. A., Kramer, M. R., Waller, L. A. and Martin, G. S. (2018) Risk Factors for Septicemia Deaths and Disparities in a Longitudinal US Cohort, *Open Forum Infectious Diseases*, 5(12), ofy305.

19. Mayr, F. B., Yende, S., Linde-Zwirble, W. T., Peck-Palmer, O. M., Barnato, A. E., Weissfeld, L. A. and Angus, D. C. (2010) Infection rate and acute organ dysfunction risk as explanations for racial differences in severe sepsis, *JAMA*, 303(24), 2495-503.

20. Melamed, A., & Sorvillo, F. J. (2009). The burden of sepsis-associated mortality in the United States from 1999 to 2005: an analysis of multiple-cause-of-death data. *Critical Care*, 13(1), R28.

21. Matter, M. L., Shvetsov, Y. B., Dugay, C., Haiman, C. A., Le Marchand, L., Wilkens, L. R., & Maskarinec, G. (2017). High mortality due to sepsis in Native Hawaiians and African Americans: the multiethnic cohort. *PLoS one*, 12(5), e0178374.

22. Schrader, C. D. and Lewis, L. M. (2013) Racial disparity in emergency department triage, *J Emerg Med*, 44(2), 511-518.

23. Raman, J., Johnson, T. J., Hayes, K. and Balamuth, F. (2019) Racial Differences in Sepsis Recognition in the Emergency Department, *Pediatrics*, 144(4), e20190348.
24. Alanee, S., Holland, B., Clemons, J. and Dynda, D. (2018) Death due to sepsis in patients diagnosed with prostate cancer, *The Prostate*, 79(3), 295-301.
25. Nafiu, O. O., Mpodoy, C., Kim, S. S., Uffman, J. C. and Tobias, J. D. (2020) Race, Postoperative Complications, and Death in Apparently Healthy Children, *Pediatrics*, 146(2).
26. Bauer, M. E., Bateman, B. T., Bauer, S. T., Shanks, A. M. and Mhyre, J. M. (2013) Maternal sepsis mortality and morbidity during hospitalization for delivery: temporal trends and independent associations for severe sepsis, *Anesthesia & Analgesia*, 117(4), 944-50.
27. Bakalli, H., Close, R., & Kellywood, K. (2020). SEPSIS IN THE NATIVE AMERICAN POPULATION. *Chest*, 158(4), A707.
28. Jones, J. M., Fingar, K. R., Miller, M. A., Coffey, R., Barrett, M., Flottemesch, T., Heslin, K. C., Gray, D. T. and Moy, E. (2017) Racial Disparities in Sepsis-Related In-Hospital Mortality: Using a Broad Case Capture Method and Multivariate Controls for Clinical and Hospital Variables, 2004-2013, *Critical Care Medicine*, 45(12), e1209-e1217.
29. Jacobs, Z. G., Prasad, P. A., Fang, M. C., Abe-Jones, Y. and Kangelaris, K. N. (2019) The Association between Limited English Proficiency and Sepsis Mortality, *Journal of Hospital Medicine*, 14, E1-E7.
30. Thavamani, A., Umapathi, K. K., Dhanpalreddy, H., Khatana, J., Chotikanatis, K., Allareddy, V. and Roy, A. (2020) Epidemiology, Clinical and Microbiologic Profile and Risk Factors for Inpatient Mortality in Pediatric Severe Sepsis in the United States From 2003 to 2014: A Large Population Analysis, *Pediatric Infectious Disease Journal*, 39(9), 781-788.
31. Chang, D. W., Tseng, C. H., and Shapiro, M. F. (2015). Rehospitalizations following sepsis: common and costly. *Critical Care Medicine*, 43(10), 2085.
32. Kumar, G., Taneja, A., Majumdar, T., Jacobs, E. R., Whittle, J., Nanchal, R. (2014) The association of lacking insurance with outcomes of severe sepsis: retrospective analysis of an administrative database, *Critical Care Medicine*, 42(3), 583-91.
33. Bohanon, F. J., Nunez Lopez, O., Adhikari, D., Mehta, H. B., Rojas-Khalil, Y., Bowen-Jallow, K. A. and Radhakrishnan, R. S. (2017) Race, Income, and Insurance Status

Affect Neonatal Sepsis Mortality and Healthcare Resource Utilization, *Pediatric Infectious Disease Journal*, 37(7), e178.

34. Odetola, F. O., & Gebremariam, A. (2019). Resource Use and Outcomes for Children Hospitalized With Severe Sepsis or Septic Shock. *Journal of Intensive Care Medicine*, 36(1), 89-100.

35. Kanter, G. P., Segal, A. G. and Groeneveld, P. W. (2020) Income Disparities In Access To Critical Care Services, *Health Affairs*, 39(8), 1362-1367.

36. Barbash, I. J. (2020) Disparities in Sepsis Outcomes: A Problem in Need of Solutions, *Critical Care Medicine*, 48(7), 1079-1080.

37. Rush, B., Danziger, J., Walley, K. R., Kumar, A. and Celi, L. A. (2020) Treatment in Disproportionately Minority Hospitals Is Associated With Increased Risk of Mortality in Sepsis: A National Analysis, *Critical Care Medicine*, 48(7), 962-967.

38. Corl, K., Levy, M., Phillips, G., Terry, K., Friedrich, M. and Trivedi, A. N. (2019) Racial And Ethnic Disparities In Care Following The New York State Sepsis Initiative, *Health Affairs*, 38(7), 1119-1126.

39. Sepsis Alliance Awareness Survey. (2020) <https://www.sepsis.org/2020-sepsis-awareness-survey/>

40. Disparities. (2019, October) <https://www.ihs.gov/newsroom/factsheets/disparities/>